

Aortic arch surgery

Background

The aorta, the largest blood vessel in the body, emerges from the heart in an upward direction before bending backwards and downwards to supply the lower parts of the body. This curvature, known as the aortic arch, also gives off branches that supply the brain and arms. When the walls of the aortic arch dilate and widen (causing an aneurysm), or when a tear appears (aortic dissection), surgical or medical intervention is required, as severe internal bleeding can occur. Risk factors for aortic aneurysms include hardening of arteries (atherosclerosis), high blood pressure, some genetic conditions (such as Marfan's Syndrome), tissue pathologies which weaken the aortic wall, or trauma.

Seeking help

Aneurysms along the aortic arch often develop slowly over many years with little or no symptoms, and are usually detected when medical imaging is conducted for unrelated reasons. Some patients experience chest pain, fast pulse, fatigue or shortness of breath. There may also be a marked difference in blood pressure measured in the arms and legs. Should these symptoms occur, it is important to seek urgent medical attention, particularly to rule out aortic dissection, which is life-threatening and demands immediate surgery.

Surgical management

Surgery is required when the aneurysm has reached a certain size, when it is ruptured, or when aortic dissection is detected.

Aortic arch surgery is a highly complex operation. It requires heart-lung bypass, whereby after opening the chest, circulation is diverted from the heart and lungs and oxygenated blood is pumped around the body using an external machine. The heart then stops beating. Subsequently, the body is cooled down to around 18–28 °C (64–82 °F), which reduces the metabolic demands of vital organs, allowing general circulation around the body to be stopped (circulatory arrest). Concurrently, blood is selectively pumped to the brain via arteries to maintain neurological function.

Operation on the aortic arch is then performed, which is usually replacement of diseased portions with synthetic material. Once all the blood vessels are satisfactorily replaced and connected, the blood is slowly warmed and introduced into the body, and the heart restarted. During this operation, repairs can also be performed for any concomitant heart anomalies, such as blocked coronary arteries.

Depending on the nature of the aortic aneurysm, an endovascular approach, whereby a metal wire mesh (stent)

is inserted via an artery to stabilise the aneurysm, can be utilised. This minimally-invasive procedure removes the need for open heart surgery in some select cases.

Medical management

Adequate blood pressure control is a necessity in all patients. Anti-cholesterol drugs, such as statins, may also be given to lower cholesterol levels and control inflammation. Long term check-ups are required.

Risks

The complexity of aortic arch surgery carries with it serious risk. Complications include:

- Stroke
- Damage to spinal cord
- Bleeding
- Heart attacks
- Lung and/or kidney failure
- Bowel necrosis
- Risk to life

However, despite these risks, long term outcome is reasonable, although this is dependent upon the type of arch disorder. Significantly improved quality of life usually follows surgery. Rates of risk vary between institutions, as well as with patient comorbidities and anatomy. Such risks, balanced against potential benefits, should be discussed with your surgeon.

For more information, please visit the following websites:
<http://emedicine.medscape.com/article/424904-overview>
http://my.clevelandclinic.org/heart/disorders/aorta_marfan/complexaortasurgery.aspx
<http://www.sts.org/patient-information/aneurysm-surgery/aortic-aneurysms>

doi: 10.3978/j.issn.2225-319X.2013.01.12

Section Writer: David H. Tian

Illustration Editor: Beth Croce

Section Editor: Ashutosh Hardikar

For specific information concerning your medical condition, ACS suggests that you consult your physician. This page may be photocopied noncommercially by physicians to share with patients.

